# Human Fc gamma RIIa / CD32a (167R) protein

Catalog number HUGR2AR-U



#### **Synonyms**

CD32, CD32A, FCGR2, FCGR2A, FCGRII, FCGRIIA, FCR2, FCR2A, FCRII, FCRIIA, IGFR2, IGFR2A, IGFRII, IGFRIIA, Fc gamma R2, Fc gamma R1, Fc gamma RIIa, Fcg R2, Fcg R2a, Fcg RII, Fcg RIIa

# **Species**

Human

### **Accession number**

P12318

# **Allotype**

167R

# **Conjugation status**

Unconjugated (no label). The protein contains an AVI tag but this has not been biotinlyated in this product.

# **Purity**

>95% monomer purity as determined by SDS-PAGE and SEC-HPLC.

#### **Endotoxin**

<1.0 EU per mg as determined by the LAL method.

### **Protein design**

The sequence of the extracellular domain of human CD32a (Ala 36-Ile 218) was fused with a C-terminal tag consisting of the AVI tag, TEV protease recognition sequence and a 10-His tag. The full protein sequence can be downloaded from the product webpage.

### Molecular weight

The recombinant human CD32a (167R) including tag consists of 222 amino acids and has a theoretical mass of 24792 Da.

### **Expression host**

Human embryonic kidney (HEK) 293 cells.

# **Formulation**

Lyophilized from sterile PBS, pH 7.4. No preservatives or cryoprotectants have been added.

# Reconstitution

To obtain a final concentration of 1 mg/ml reconstitute 250  $\mu$ g vials with 250  $\mu$ l water and 1.0 mg vials with 1.0 ml water.

Solubilize for 30 to 60 minutes at room temperature with occasional gentle mixing. Do not vortex.

# **Shipping**

All recombinant proteins are provided as lypholized powder and shipped at ambient temperature.

# Storage and stability

Lyophilized proteins are stable at ambient temperature for at least 2 weeks. If the protein is not to be used immediately then the protein should be stored in lyophilized form at -20 °C for up 12 months. Once the protein has been reconstituted we recommend storage at 4 °C for up to one week. For longer term storage of protein in solution we recommend aliquoting into smaller vials to avoid repeated freeze-thaw cycles and storage at -20 or -80 °C for up to 3 months.

# **Quality control**

All recombinant proteins are tested for purity by SDS-PAGE and SEC-HPLC with a minimum requirement of >95% monomer purity. Biological activity is confirmed by surface plasmon resonance on a Biacore instrument. Please see certificate of analysis (COA) for batch specific quality control data and images.

# **Product description**

Low affinity immunoglobulin gamma Fc receptor IIa, also known as FcyRIIa or CD32a, is a type I integral membrane glycoprotein. CD32a is a member of the immunoglobulin superfamily and is expressed on macrophages, monocytes, neutrophils, eosinophils and dendritic cells, epithelial cells, platelets and activated CD4+ T cells. CD32a binds monomeric IgG with low affinity but is very efficient at binding immune complexes and is involved in phagocytosis and clearing of immune complexes. CD32a is structurally composed of two extracellular immunoglobulin domains of the C2-type that interact with the IgG Fc domain, a transmembrane domain and a short cytoplasmic tail containing the immunoreceptor tyrosine-based activation (ITAM) motif. The product provided only contains the extracellular portion of CD32a.

CD32a has two allotypic variants differing at amino acid position 167, one containing histidine (H167) and the other arginine (R167). H131 exhibits a higher affinity to human lgG1 and lgG2 than the R167 does and is thought to be primarily responsible for the phagocytosis of lgG-opsonized bacteria.