

Mouse Fc gamma RIIIa / CD16a protein

Catalog number MOGR3A-U

Synonyms

CD16, CD16A, FCGR3, FCGR3A, FCGRIII, FCGRIIIa, FCR3, FCR3A, FCRIII, FCRIIIa, IGFR3, IGFR3A, IGFRIII, IGFRIIIa, Fc gamma R3, Fc gamma R3a, Fc gamma RIII, Fc gamma RIIIa, Fcg R3, Fcg R3a, Fcg RIII, Fcg RIIIa

Species

Mouse

Accession number

P08508

Allotype

Not applicable

Conjugation status

Unconjugated

Purity

>95% monomer purity as determined by SDS-PAGE and SEC-HPLC.

Endotoxin

<1.0 EU per mg as determined by the LAL method.

Protein design

The sequence of the extracellular domain of mouse CD16a (Leu 32-Thr 215) was fused with a C-terminal tag consisting of the AVI tag, TEV protease recognition sequence and a 10-His tag. The full protein sequence can be downloaded from the product webpage.

Molecular weight

The recombinant mouse CD16a including tag consists of 223 amino acids and has a theoretical mass of 25543 Da.

Expression host

Human embryonic kidney (HEK) 293 cells

Formulation

Lyophilized from sterile PBS, pH 7.4. No preservatives or cryoprotectants have been added.

Reconstitution

To obtain a final concentration of 1 mg/ml reconstitute 250 µg vials with 250 µl water and 1.0 mg vials with 1.0 ml water.

Solubilize for 30 to 60 minutes at room temperature with occasional gentle mixing. Do not vortex.

Shipping

All recombinant proteins are provided as lyophilized powder and shipped at ambient temperature.

Storage and stability

Lyophilized proteins are stable at ambient temperature for at least 2 weeks. If the protein is not to be used immediately then the protein should be stored in lyophilized form at -20 °C for up to 12 months. Once the protein has been reconstituted we recommend storage at 4 °C for up to one week. For longer term storage of protein in solution we recommend aliquoting into smaller vials to avoid repeated freeze-thaw cycles and storage at -20 or -80 °C for up to 3 months.

Quality control

All recombinant proteins are tested for purity by SDS-PAGE and SEC-HPLC with a minimum requirement of >95% monomer purity. Biological activity is confirmed by surface plasmon resonance on a Biacore instrument. Please see certificate of analysis (COA) for batch specific quality control data and images.

Product description

Low affinity immunoglobulin gamma Fc receptor IIIa, also known as FcγRIIIa or CD16a, is a type I integral membrane glycoprotein. CD16a is a member of the immunoglobulin superfamily and is expressed on macrophages, monocytes and NK cells. CD16a binds monomeric IgG with low affinity but is efficient at binding immune complexes and functions in NK cell activation, phagocytosis and antibody-dependent cellular cytotoxicity (ADCC). CD16a is structurally composed of two extracellular immunoglobulin domains of the C2-type that interact with the IgG Fc domain, a transmembrane domain and a short cytoplasmic tail. CD16a is associated with a dimer of the common Fc receptor gamma-chain which contains the immunoreceptor tyrosine-based activation (ITAM) motif. The product provided only contains the extracellular portion of CD16a.

For research use only. Not for use in diagnostic or therapeutic procedures.

Gamma Proteins Ltd, Portland House, Belmont Business Park, Durham, DH1 1TW, United Kingdom
Email: support@gammaproteins.com Website: www.gammaproteins.com