

Cynomolgus FcRn (FCGRT-B2M) heterodimer protein

Catalog number CYFCRN-U

Synonyms

FcRn, FCGRT, FCGRT & B2M, FCGRT and B2M, Neonatal Fc receptor, Neonatal receptor, Brambell receptor

Species

Cynomolgus

Accession number

Q8SPV9 / Q8SPW0

Allotype

Not applicable

Conjugation status

Unconjugated (no label). The protein contains an AVI tag but this has not been biotinylated in this product.

Purity

>95% monomer purity as determined by SDS-PAGE and SEC-HPLC.

Endotoxin

<1.0 EU per mg as determined by the LAL method.

Protein design

The sequence of the extracellular domain of cynomolgus FCGRT (Ala 24-Ser 297) was fused with a C-terminal tag consisting of the AVI tag, TEV protease recognition sequence and a 10-His tag. This was co-transfected with the sequence of cynomolgus B2M (Ile 21-Met 119) and the resulting FcRn heterodimer was purified. The full protein sequences can be downloaded from the product webpage.

Molecular weight

The recombinant FcRn heterodimer including tags consists of 412 amino acids and has a theoretical mass of 46541 Da. The heterodimer consists of cynomolgus FCGRT (313 amino acids; 34898 Da) and cynomolgus B2M (99 amino acids; 11643 Da).

Expression host

Human embryonic kidney (HEK) 293 cells.

Formulation

Lyophilized from sterile PBS, pH 7.4. No preservatives or cryoprotectants have been added.

Reconstitution

To obtain a final concentration of 1 mg/ml reconstitute 250 µg vials with 250 µl water and 1.0 mg vials with 1.0 ml water. Solubilize for 30 to 60 minutes at room temperature with occasional gentle mixing. Do not vortex.

Shipping

All recombinant proteins are provided as lyophilized powder and shipped at ambient temperature.

Storage and stability

For long term storage the protein should be stored in lyophilized form at -20 °C for up to 12 months from date of receipt. Lyophilized proteins are stable at ambient temperature for at least 2 weeks.

Once the protein has been reconstituted we recommend storage at 4 °C for up to one week. For longer term storage of protein in solution we recommend aliquoting into smaller vials to avoid repeated freeze-thaw cycles and storage at -20 or -80 °C for up to 12 months.

Quality control

All recombinant proteins are tested for purity by SDS-PAGE and SEC-HPLC with a minimum requirement of >95% monomer purity. Biological activity is confirmed by surface plasmon resonance on a Biacore instrument. Please see certificate of analysis (COA) for batch specific quality control data and images.

Product description

IgG Fc fragment receptor transporter, also known as the neonatal Fc receptor or more commonly FcRn, consist of two subunits (FCGRT and B2M) and forms an MHC class I-like heterodimer. FcRn binds to the Fc domain of monomeric IgG and mediates the pH dependent recycling of IgG as well as its uptake from milk and transfer from mother to fetus.

For research use only. Not for use in diagnostic or therapeutic procedures.

Gamma Proteins Ltd, Portland House, Belmont Business Park, Durham, DH1 1TW, United Kingdom
Email: support@gammaproteins.com Website: www.gammaproteins.com